


« Greens for Nuclear Energy      The declining value of wind and solar electricity »

Radiation Amplification Factor RAF in April 2021

We had a period of several cloudless, blue sky days at the end of April 2021. So time to redo a calculation of the Radiation Amplification Factor RAF. In short, we want to see how the variation of the Total Ozone Column (TOC) influences the effective UVB radiation at ground level. I wrote several time on this, and usually we found an RAF of approx 1.05 to 1.10.

First here a graph showing the variation of total solar irradiance (blue curve, unit W/m2) and the effective UVB (red curve, unit mMED/h):



First remark that the peak solar irradiance was practically constant; the 24th April was a bit hazy, so it will be left out in the computations. The numbers in the turquoise boxes are the maximum values of the TOC, measured in DU (Dobson Unit) with our Microtops II instrument (serial 5375). Let us first plot the UVBeff versus the TOC:




fig. 1 UVBeff versus maximum daily TOC (5 days: 23 and 25 to 28 April 2021)

Clearly the UVBeff values decrease with increasing TOC, as the thicker ozone column filters out more UVB radiation. The empirical relationship is practically linear, and suggests that a dip of 100 DU (a quite substantial thinning of the ozone layer) would cause an increase of effective UVB of about 0.6 MED/h or 1.7 UVI (as 1 MED/h = 25/9 UVI).


The numerical correct definition of the RAF is :  $UVB = C * TOC^{RAF}$  where \*\* means "at the power of" Taking the natural logarithm gives  $\ln(UVB) = \ln(C) + RAF * \ln(TOC)$  or  $RAF = [\ln(UVB) - \ln(C)] / \ln(TOC)$ .

If we have many measurement couples of UVB and TOC, it can be shown (see [here](#)) that

$RAF = [-\ln(UVB_i / UVB_0)] / [\ln(TOC_i / TOC_0)]$

where the index i corresponds to the ith measurement couple, and 0 to that taken as a reference (usually i=0). This is equivalent to say that RAF is the slope of the linear regression line through the scatterplot of  $-1 * \ln(UVB_i / UVB_0)$  versus  $\ln(TOC_i / TOC_0)$ .

Here is that plot:




RAF computed from TOC

The slope is 1.0461, so the (erythemal) RAF computed from the 5 blue sky days is  $RAF = 1.0461 \sim 1.05$


This has to be compared to the value  $RAF = 1.08$  in the referenced paper [ref. 1]. Note the excellent  $R^2 = 0.96$  of this linear fit.

There is some discussion if TOC should be replaced by  $TOC_{slant} = TOC / \cos(SZA)$ , where SZA is the solar zenith angle. If we do this, the  $RAF \sim 1.10$ , close to the previous value; the  $R^2$  is somewhat lower with  $R^2=0.91$ . The SZA is practically constant for the 5 days with  $SZA \sim 38^\circ$ .



RAF computed from TOC slant =  $TOC / \cos(SZA)$

The  $RAF = 1.10$  value is close to what Jay Herman published in GRL in figure 8 [ref. 2] (red lines added):



RAF from Erythemal UVB as a function of SZA

Conclusion

These 5 days of cloudless sky give practically the same results for RAF as that found during previous investigations. As a very raw rule of thumb one could keep in mind that a dip of 100 DU yields an increase of at most 2 UVI. The following table resumes the findings of this paper and the references 1 to 5:

reference	year	author	RAF (TOC not slanted)	Month
this paper	2021	Massen	1.05	April
1	2013	Massen	1.08	April
2	2010	Herman	1.70	
3	2014	Massen	1.17	March
4	2016	Massen	1.10	April
5	2018	Massen	1.09	September

Table of erthymal RAF's

References:

- [1] MASSEN, Francis, 2013: Computing the Radiation Amplification Factor RAF using a sudden dip in Total Ozone Column measured at Diekirch, Luxembourg ([link](#))
- [2] HERMAN, Jay, 2010: Use of an improved radiation amplification factor to estimate the effect of an ozone changes on action spectrum weighted irradiances and an instrument response function. Journal of Geophysical Research, vol.115, 2010 ([link](#))
- [3] MASSEN, Francis, 2014 : RAF revisited ([link](#))
- [4] MASSEN, Francis, 2016: First Radiation Amplification Factor for 2016 ([link](#))
- [5] MASSEN, Francis, 2018: UVI and Total Ozone ([link](#))

Share this:

Facebook X

Loading...

Related

UVI and total ozone column: 3 days in September 2018  
September 27, 2018  
With 2 comments

RAF in Diekirch, LU, March 2022 | meteoLCD Weblog Says:  
March 13, 2022 at 10:45 | Reply

First Radiation Amplification Factor for April 21, 2016  
With 4 comments

This entry was posted on April 29, 2021 at 14:42 and is filed under Uncategorized. You can follow any responses to this entry through the [RSS 2.0](#) feed. You can [leave a response](#), or [trackback](#) from your own site.

One Response to “Radiation Amplification Factor RAF in April 2021”

- RAF in Diekirch, LU, March 2022 | meteoLCD Weblog Says: March 13, 2022 at 10:45 | Reply
- [...] [1] MASSEN, Francis: Radiation Amplification Factor in April 2021 (link)
- [...]

Leave a comment